

Hair-Related Dysmorphophobia: A Mini-Review

Ademir Carvalho Leite Júnior¹

¹Scientific Director at HTRI Clinic

Correspondence: ademirjr@ademirjr.com.br

Abstract

Hair-related dysmorphophobia, more accurately termed hair-related body dysmorphic disorder (BDD), is a psychiatric condition characterized by obsessive preoccupation with perceived defects in hair appearance that are not observable or appear minor to others, causing significant distress and functional impairment. This mini-review examines clinical characteristics, prevalence, manifestations, relationship with other disorders, diagnostic criteria, therapeutic approaches, and psychosocial impacts of this condition. Through analysis of current literature, we observe that 29.6% of patients seeking treatment for hair loss complaints meet criteria for BDD, representing approximately 10 times higher prevalence than in general dermatology patients (2.7%)¹. Early recognition of this condition is fundamental to avoid unnecessary treatments and promote appropriate psychotherapeutic and pharmacological interventions. Assessment should include standardized anxiety instruments and specific BDD scales, allowing for a multidisciplinary approach that integrates dermatological and psychiatric aspects. The condition significantly impacts quality of life, with high rates of depression, anxiety, and social impairment. Treatment requires an interdisciplinary approach combining cognitive-behavioral therapy and selective serotonin reuptake inhibitors as first-line interventions. Future research should focus on developing hair-specific assessment and treatment protocols, large-scale epidemiological studies in diverse populations, and investigation of cultural and social factors influencing hair-related appearance standards.

Keywords: Body dysmorphic disorder, Hair-Related dysmorphophobia, Hair Loss, anxiety, psychotrichology

Citation: Leite Jr AC. Hair-Related Dysmorphophobia: A Mini-Review. Brazilian Journal of Hair Health. 2025;2:bjhh35.

doi: <https://doi.org/10.62742/2965-7911.2025.2.bjhh35>

Received: July 30, 2025

Revised: August 08, 2025

Accepted: August 13, 2025

Published: August 14, 2025



Copyright: This content is licensed under the terms and conditions of the Creative Commons Attribution 4.0 International License (CC BY).

1. Introduction

The relationship between psychological aspects and hair manifestations has been the subject of growing scientific interest in recent decades. Hair-related dysmorphophobia, currently termed hair-related body dysmorphic disorder (BDD), represents a specific manifestation of a broader psychiatric disorder that significantly affects the quality of life of affected individuals^{1, 2}.

Hair-related BDD is classified within the obsessive-compulsive and related disorders (OCRD) category in DSM-5, representing a significant clinical challenge due to its impact on psychological well-being and social functioning¹. BDD is defined as a psychiatric condition characterized by disabling preoccupation with non-existent or slight defects in appearance that are not observable or appear only slight to others¹. When focused on hair, this manifests as excessive concern with perceived hair defects, leading to compulsive behaviors and significant functional impairment³.

Understanding this condition requires a multidisciplinary approach that integrates knowledge from dermatology, psychiatry, and psychology, recognizing that hair possesses profound symbolic meanings related to identity, femininity, masculinity, and self-image. As demonstrated in previous studies, hair can represent fundamental elements of personal identity, becoming a target for dysmorphic concerns in predisposed individuals.

The interface between psychology and hair manifestations, termed psychotrichology, offers important perspectives for understanding hair-related BDD. Stressful life events can trigger or exacerbate hair problems through complex neuroimmunoendocrine mechanisms, suggesting that psychological factors play a crucial role in hair health and related disorders.

Clinical Definition and Characteristics

Hair-related BDD presents specific clinical features that distinguish it from other normal appearance concerns. The condition manifests through a complex interplay of cognitive, behavioral, and physical symptoms that create significant distress and functional impairment^{1, 2, 3}. Obsessive preoccupation with hair appearance, texture, thickness, or pattern constitutes the central core of the condition, with patients developing intrusive thoughts about hair appearance that persist despite attempts to dismiss them. These cognitive distortions often involve catastrophic interpretations of minor hair changes, excessive self-monitoring and comparisons with others, and persistent beliefs that they are being negatively judged due to their hair appearance¹.

The behavioral manifestations of hair-related BDD represent attempts to alleviate the intense anxiety generated by dysmorphic concerns. Patients typically engage in repetitive mirror checking or, conversely, complete mirror avoidance when the anxiety becomes overwhelming. Excessive hair grooming, manipulation, and styling behaviors are common, often accompanied by frequent hair washing or compulsive touching of the hair. Many individuals seek constant reassurance from others about their hair appearance, yet remain unconvinced by positive feedback. Social avoidance becomes increasingly prominent, with patients withdrawing from situations where they perceive their hair might be scrutinized. Camouflaging behaviors, such as wearing hats, choosing specific hairstyles, or positioning themselves to hide perceived defects, become integral parts of their daily routine¹.

Paradoxically, individuals with hair-related BDD may engage in self-destructive behaviors that effectively damage the hair they are so concerned about protecting. This creates a vicious cycle where real hair problems develop secondary to the psychological condition, reinforcing the patient's distorted beliefs and intensifying their distress². The complexity of this phenomenon highlights the sophisticated interplay between psychological distress and physical manifestations, emphasizing the critical need for early recognition and intervention to prevent the escalation of both psychological and dermatological complications.

Prevalence and Demographics

Epidemiological data reveal a surprisingly high prevalence of BDD among patients seeking treatment for hair issues, fundamentally challenging our understanding of the relationship between dermatological complaints and underlying psychiatric conditions. The landmark study by Doğruk Kaçar et al.¹ demonstrated that 29.6% of patients presenting hair loss complaints meet criteria for BDD, representing approximately 10 times higher prevalence than in general dermatology patients (2.7%)¹. This striking disparity suggests that hair-related concerns may be particularly vulnerable to dysmorphic interpretation, possibly due to the high visibility and cultural significance of hair in personal identity and social perception.

The gender distribution of hair-related BDD presents interesting patterns that contrast with general BDD prevalence. Males demonstrate significantly higher rates of BDD among hair loss patients (52.4%) compared to females (25.6%)¹, a finding that diverges from the general prevalence of BDD in the population, where no significant gender differences are typically observed⁴. This gender disparity in hair-related presentations may reflect cultural and social factors that place particular emphasis on male hair as a symbol of youth, virility, and attractiveness. The psychological impact of hair loss on male identity appears to create unique vulnerabilities to dysmorphic interpretations that warrant specific clinical attention.

In the general population, BDD prevalence ranges from 0.5% to 3.2%, with relatively equal distribution across genders⁴. However, when examining specific body areas of concern, hair-related preoccupations show distinct demographic patterns that suggest hair holds particular significance for certain populations. The elevated prevalence among males seeking hair loss treatment indicates that this demographic represents a high-risk group requiring targeted screening and intervention strategies. These epidemiological findings have important implications for clinical practice, suggesting that routine psychiatric screening should be considered standard care in dermatological settings treating hair-related complaints.

Relationship with Body Dysmorphic Disorder and Trichotillomania

Hair-related BDD exists within a complex relationship with other hair-focused disorders, creating diagnostic and therapeutic challenges that require careful clinical consideration. The relationship with trichotillomania illustrates the complexity of hair-related psychopathology, as both conditions involve significant distress related to hair but differ fundamentally in their manifestations and underlying mechanisms. Trichotillomania involves the compulsive act of hair pulling, resulting in noticeable hair loss and often leaving distinctive patterns of alopecia^{5,6,7}. In contrast, hair-related BDD focuses on perceived defects in hair appearance without necessarily involving hair-pulling behaviors, though the distress and functional impairment can be equally severe¹.

The potential for comorbidity between these conditions adds another layer of complexity to clinical assessment and treatment planning. Some individuals experience both hair-pulling urges characteristic of trichotillomania and dysmorphic concerns about hair appearance typical of BDD⁸. This overlap highlights the intricate nature of hair-related disorders and emphasizes the need for comprehensive evaluation that considers multiple diagnostic possibilities. Clinicians must be prepared to identify and address both conditions when they co-occur, as treatment approaches may need to be modified to address the full spectrum of symptoms and behaviors.

Within the broader spectrum of BDD, hair concerns represent one of the most common areas of focus, alongside skin, nose, and other facial features². The underlying psychological mechanisms remain remarkably consistent across different body areas of concern, involving similar patterns of perfectionism, cognitive distortions, and appearance-related anxiety. However, hair-related BDD may have unique features related to the particular symbolic and cultural significance of hair. The visibility of hair, its role in personal identity, and its association with attractiveness, youth, and social status create specific vulnerabilities that may require targeted therapeutic approaches. Understanding these relationships is crucial for developing effective treatment strategies that address not only the symptoms but also the underlying psychological mechanisms that drive the condition.

Diagnostic Criteria and Assessment

The diagnostic framework for hair-related BDD follows the established DSM-5 criteria for body dysmorphic disorder, adapted specifically for hair-related concerns. The core diagnostic criteria include preoccupation with one or more perceived defects in hair appearance that are not observable or appear minor to others, representing the fundamental disconnect between subjective perception and objective reality that characterizes this condition. Patients must demonstrate repetitive behaviors or mental acts in response to their appearance concerns, such as mirror checking, grooming rituals, or seeking reassurance. The condition must cause clinically significant distress or impairment in social, occupational, or other important areas of functioning, distinguishing pathological preoccupation from normal appearance concerns. Additionally, the symptoms cannot be better explained by concerns about body weight in individuals with eating disorders².

The assessment of hair-related BDD requires a multimodal approach that combines validated instruments with clinical interview techniques. The Body Dysmorphic Disorder Questionnaire (BDDQ) has proven particularly valuable, as demonstrated in the landmark study showing 29.6% prevalence among hair loss patients¹. The Yale-Brown Obsessive-Compulsive Scale Modified for BDD (BDD-YBOCS) provides a standardized measure of symptom severity, allowing clinicians to track treatment response and monitor disease progression⁹. Clinical interviews focusing on appearance-related distress and functional impairment remain fundamental for comprehensive assessment, providing context and nuance that standardized instruments may miss.

The assessment of anxiety represents a crucial component of evaluation, given the high level of anxiety consistently associated with hair-related BDD. Multiple studies demonstrate that anxiety is frequently present in patients with hair loss^{10,11}, making standardized anxiety assessment tools essential for comprehensive care. The Beck Anxiety Inventory (BAI) and Hamilton Anxiety Rating Scale (HAM-A) are widely used instruments that complement specific BDD assessment and provide valuable information about comorbid anxiety symptoms. The Hospital Anxiety and Depression Scale (HADS) has also shown utility in dermatological settings, including studies with hair-related conditions¹². These anxiety assessment tools not

only aid in initial diagnosis but also provide valuable monitoring capabilities during treatment, helping clinicians adjust therapeutic approaches based on symptom response and overall clinical improvement.

Therapeutic Approaches and Effectiveness

The treatment of hair-related BDD requires a comprehensive approach that addresses both the psychological and dermatological aspects of the condition. Cognitive Behavioral Therapy (CBT) stands as the first-line psychological treatment for BDD, including hair-related presentations, demonstrating consistent efficacy across multiple studies and clinical populations. The therapeutic framework of CBT incorporates several key components that directly target the core symptoms of hair-related BDD. Cognitive restructuring techniques help patients identify and challenge distorted thoughts about their hair appearance, addressing the catastrophic interpretations and negative self-evaluations that maintain the disorder. Exposure and response prevention protocols systematically reduce avoidance behaviors while helping patients tolerate the anxiety associated with hair-related situations. Behavioral experiments allow patients to test their catastrophic predictions about how others perceive their hair, often revealing the discrepancy between feared and actual outcomes².

Emerging therapeutic approaches show promise for enhancing treatment outcomes in hair-related BDD. Acceptance and Commitment Therapy (ACT) represents a newer intervention that focuses on psychological flexibility and value-based living rather than symptom elimination. While research specifically examining ACT for hair-related BDD presentations remains limited, preliminary studies suggest potential benefits for BDD and related disorders¹³. The integration of mindfulness-based approaches may also provide valuable tools for managing the intense anxiety and rumination characteristic of hair-related BDD.

Pharmacological interventions play a crucial role in the comprehensive treatment of hair-related BDD, with Selective Serotonin Reuptake Inhibitors (SSRIs) representing the gold standard for medication management. Clinical experience indicates that higher doses are often required compared to depression treatment, reflecting the neurobiological differences underlying BDD. Treatment duration typically extends 12-16 weeks for optimal response, with many patients requiring longer-term maintenance therapy to prevent relapse⁴. The medication management requires careful monitoring and adjustment, as individuals with BDD may be particularly sensitive to side effects that could affect their appearance concerns.

Alternative pharmacological approaches may be considered for treatment-resistant cases or as augmentation strategies. N-acetylcysteine has shown promise for glutamate system modulation in obsessive-compulsive and related disorders¹⁴. Tricyclic antidepressants may be beneficial in cases that do not respond adequately to SSRIs, while antipsychotic augmentation can be considered in severe cases where delusional thinking predominates¹⁵. The selection of pharmacological interventions must be individualized based on symptom severity, comorbid conditions, and patient preferences.

The complex nature of hair-related BDD necessitates interdisciplinary treatment approaches that integrate expertise from multiple specialties. Collaboration between dermatologists, psychiatrists, and psychologists provides the most comprehensive care, addressing both the psychological aspects of the condition and any underlying dermatological concerns². This integrated approach recognizes that many patients with hair-related BDD may have concurrent dermatological conditions that require medical management alongside psychological treatment. The coordination between specialists ensures that all aspects of the patient's presentation are addressed while avoiding conflicting or counterproductive interventions.

Psychological and Social Impacts

Hair-related BDD creates profound and far-reaching consequences that extend well beyond the immediate symptoms, fundamentally altering the individual's psychological well-being and social functioning. The psychological impact manifests through multiple interconnected pathways that create a complex web of distress and impairment. High rates of depression and anxiety represent common comorbidities¹⁰, often developing as secondary consequences of the chronic distress and social isolation associated with hair-related BDD. The persistent preoccupation with perceived hair defects creates a state of chronic psychological tension that can evolve into clinically significant mood disorders, requiring additional therapeutic attention and potentially complicating treatment approaches.

The risk of suicidal ideation and behavior appears elevated in individuals with hair-related BDD compared to the general population, reflecting the severe psychological distress and hopelessness that can accompany this condition. The combination of persistent appearance-related distress, social isolation, and functional impairment creates a particularly vulnerable psychological state that requires careful monitoring and intervention. Quality of life becomes significantly compromised across multiple domains, with individuals reporting decreased satisfaction with personal relationships, professional accomplishments, and overall life fulfillment. The development of avoidant behaviors and progressive social isolation represents a particularly insidious aspect of the condition, as patients gradually withdraw from social interactions to avoid perceived scrutiny of their hair appearance¹⁶.

The social and occupational consequences of hair-related BDD create additional layers of impairment that can perpetuate and intensify the underlying psychological distress. Avoidance of social situations becomes increasingly prominent, particularly in contexts where individuals perceive their hair might be subject to observation or judgment by others. Professional and academic performance often suffers as preoccupation with hair appearance interferes with concentration, productivity, and interpersonal effectiveness. The time consumed by hair-related rituals, mirror checking, and grooming behaviors can significantly impact daily functioning and professional responsibilities.

Relationship difficulties emerge as a natural consequence of the anxiety, avoidance, and preoccupation characteristic of hair-related BDD. Intimate relationships may be strained by the individual's need for constant reassurance, avoidance of physical intimacy, or withdrawal from social activities. Family relationships can be affected by the patient's distress, time-consuming rituals, and resistance to social engagement. The economic burden associated with excessive grooming products, frequent dermatological consultations, unnecessary treatments, and cosmetic procedures creates additional stress for both patients and their families¹⁷. This financial impact often compounds the psychological distress and can create additional relationship strain, particularly when family members disagree about the necessity or appropriateness of appearance-related expenditures.

The pattern of comorbidity in hair-related BDD reveals the complex interplay between different psychiatric conditions and emphasizes the need for comprehensive assessment and treatment planning. Major depressive disorder, social anxiety disorder, and obsessive-compulsive disorder frequently co-occur with hair-related BDD^{1,10}, creating diagnostic and therapeutic challenges that require careful consideration. Additionally, many patients develop other body dysmorphic concerns affecting different body areas, suggesting that hair-related BDD may represent part of a broader pattern of appearance-related distress rather than an isolated phenomenon. This comorbidity pattern has important implications for treatment planning, as interventions must address the full spectrum of symptoms and conditions rather than focusing exclusively on hair-related concerns.

Discussion

The current evidence base for hair-related dysmorphophobia demonstrates both strengths and limitations that warrant careful consideration. The landmark study by Doğruk Kaçar et al.¹ provides robust evidence for high BDD prevalence among hair loss patients, but this represents one of the few studies specifically examining hair-related presentations. Most BDD research examines the disorder broadly without differentiating between different areas of appearance concern, creating a significant gap in our understanding of this specific manifestation.

The remarkably high prevalence of BDD among patients with hair concerns (29.6%) highlights the critical importance of psychiatric screening in dermatological practice. Healthcare providers, particularly dermatologists and primary care physicians, should be aware of this high prevalence and consider BDD in patients presenting with hair concerns¹. Early identification and appropriate referral can prevent years of unnecessary medical treatments and psychological distress. The finding that males show higher rates of hair-related BDD than females contrasts with general BDD prevalence patterns, suggesting that hair concerns may have particular significance for male identity and self-image. This observation has important implications for treatment approaches and public health initiatives, particularly given the cultural and social meanings attributed to hair across different populations.

The treatment literature, while extensive for BDD generally, lacks specificity for hair-related presentations. This represents a significant gap, as hair-related concerns may require modified approaches given the visibility and social significance of hair in many cultures. The integration of anxiety assessment tools into routine evaluation is crucial, given the high comorbidity with anxiety disorders demonstrated in multiple studies^{10, 11, 12}. The psychotrichological perspective offers valuable insights, suggesting that traumatic and conflicting life events may activate psychological complexes that, when associated with symbolic meanings related to hair, can direct stress responses specifically to hair follicles. This understanding can inform both assessment and treatment approaches, potentially leading to more targeted and effective interventions.

Psychotrichological Model




Figure 1. Psychotrichological Perspectives on Psychological Disturbances Affecting Hair.

In general Several important controversies and gaps remain in our understanding of hair-related dysmorphophobia. Most BDD research examines the disorder broadly rather than focusing on specific body areas like hair, limiting our ability to develop targeted interventions. There is a notable lack of research on whether hair-focused BDD requires modified treatment approaches compared to other BDD presentations. Cultural factors represent another significant gap, with minimal research on how cultural variations in hair significance affect BDD presentation and treatment outcomes. The impact of technology, particularly social media and digital image manipulation, on hair-related BDD remains largely unexplored despite the increasing influence of these factors on appearance standards. Prevention strategies also lack sufficient research, with limited data on early intervention and prevention approaches that could reduce the development or severity of hair-related BDD.

Future research priorities should include the development of hair-specific assessment tools and treatment protocols that account for the unique aspects of hair-related concerns. Large-scale epidemiological studies in diverse populations are needed to better understand prevalence patterns across different cultural and demographic groups. Investigation of cultural and societal factors influencing hair-related appearance standards could inform culturally sensitive treatment approaches. The effectiveness of novel treatment modalities, including digital interventions that leverage technology for therapeutic purposes, requires systematic evaluation. Neuroimaging studies of hair-related appearance processing could provide insights into the neurobiological mechanisms underlying this specific manifestation of BDD.

A clinical example illustrates the practical implications of these considerations. A 25-year-old male presents to a dermatologist with concerns about "thinning hair" that is not apparent to the clinician. Despite reassurance and normal clinical findings, the patient reports spending 2-3 hours daily examining his hair, avoiding social situations, and considering hair transplant surgery. This presentation suggests hair-related BDD requiring psychiatric evaluation and treatment rather than dermatological intervention. The condition also highlights the importance of interdisciplinary care models that integrate dermatological and psychiatric expertise to provide comprehensive treatment for patients with appearance-related concerns².

Hair-related dysmorphophobia has significant implications for both clinical practice and public health that extend beyond individual patient care. The development of screening

protocols for dermatological settings, training programs for healthcare providers, and integrated care pathways could substantially improve outcomes for affected individuals. The economic burden of this condition, manifested through unnecessary treatments and lost productivity, also warrants attention from healthcare policymakers. Early recognition and appropriate treatment could reduce healthcare costs while improving patient outcomes, making this both a clinical and economic imperative. The integration of psychotrichological understanding with traditional psychiatric and dermatological approaches offers promise for developing more comprehensive and effective treatment strategies that address both the psychological and dermatological aspects of this complex condition.

Conclusion

Hair-related dysmorphophobia, or hair-related BDD, represents a significant psychiatric condition that substantially affects the quality of life of affected individuals. With a prevalence of 29.6% among patients seeking treatment for hair loss¹, this condition requires recognition and appropriate intervention by healthcare professionals.

Clinical characteristics include obsessive preoccupation with perceived defects in hair appearance, compulsive hair-related behaviors, and significant impairment in social and occupational functioning. Assessment should incorporate standardized instruments, including anxiety scales, given the high level of anxiety associated with the condition.

Effective treatment requires a multidisciplinary approach combining psychotherapeutic interventions (primarily CBT) and pharmacological treatments (primarily SSRIs), with special attention to the specific needs of patients with hair concerns. Future research should focus on developing hair-specific assessment and treatment protocols, large-scale epidemiological studies in diverse populations, and investigation of cultural and social factors influencing hair-related appearance standards.

Early recognition of this condition can prevent years of unnecessary suffering and promote more effective treatment outcomes, highlighting the importance of professional education and awareness about this specific manifestation of body dysmorphic disorder. The integration of psychotrichological perspectives with traditional psychiatric and dermatological approaches offers promise for more comprehensive understanding and treatment of this complex condition.

Funding: None.

Institutional Review Board Statement and/or Informed Consent Statement: None.

Acknowledgments: None.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Dođruk Kaçar S, Ozuğuz P, Bağcioglu E, Coskun KS, Polat S, Karaca Ş, Özbülüt Ö. Frequency of body dysmorphic disorder among patients with complaints of hair loss. *Int J Dermatol*. 2016;55(4):415-420.
2. Harth W, Blume-Peytavi U. Psychotrichology: psychosomatic aspects of hair diseases. *J Dtsch Dermatol Ges*. 2013;11(2):125-135.
3. Hair-Related Dysmorphic Disorder in a Patient with Bipolar Disorder: Case Report. *Brazilian Journal of Hair Health*. 1(1).
4. Loewen A, Blasco-Fontecilla H, Li TMH, Bella-Fernández M, Ruiz-Antorán B. Prevalence of Body Dysmorphic Disorder in the Spanish Population: Cross-Sectional Web-Based Questionnaire Study. *JMIR Form Res*. 2024;8:e46515.
5. Zhang M. Trichotillomania (Hair-Pulling Disorder). *Autism and child psychopathology series*. 2023.
6. Sheikh A, Godil A. The Psychological Impact of Trichotillomania. 2024.

7. Rosińska K, Łojewska JN, Janicka EJ, Perko A, Rosiński M, Niedźwiedzka M, Bochenek O, Koper M. From Hair-Pulling to Trichobezoars and Rapunzel Syndrome: A Comprehensive Review of Trichotillomania. *Quality in Sport*. 2024;22:54247.
8. Modi N, Chaudhary P, Mehta R, Dave K. Case Series of Trichotillomania Presenting with Three Different DSM Entities. 2021.
9. Moreno-Amador B, Piqueras JA, Rodríguez-Jiménez T, Martínez-González AE, Cervin M. Measuring symptoms of obsessive-compulsive and related disorders using a single dimensional self-report scale. *Front Psychiatry*. 2023;14:958015.
10. Malta M, Corso A. Understanding the Association Between Mental Health and Hair Loss. *Cureus*. 2025;17(1):e84777.
11. Orhan FO, Ozer A. Evaluation of anxiety and levels of serum B12, folate, TSH, ferritin, and zinc in telogen alopecia patients with trichodynia. *Int J Trichol*. 2012;4(4):251-256.
12. Christou E, Lalagianni M, McSweeney S, Cotter I, Ung CY, Walburn J, McCrone P, Turner H, McGrath JA, Weinman J, Tziotzios C. Psychosocial burden and the impact of illness perceptions and stigma on quality of life, anxiety and depression in alopecia areata: results from the Alopecia+Me study. *Br J Dermatol*. 2025;192(2):189-197.
13. Twohig MP, Capel J, Levin ME. A Review of Research on Acceptance and Commitment Therapy for Anxiety and Obsessive-Compulsive and Related Disorders. *Psychiatr Clin North Am*. 2024;47(4):729-747.
14. Parli SE, Gales MA, Gales BJ. N-Acetylcysteine for Obsessive-Compulsive and Related Disorders in Children and Adolescents: A Review. *Ann Pharmacother*. 2023;57(5):518-529.
15. Ha J, Magid M. Psychopharmacology in dermatology: Five common disorders. *Clin Dermatol*. 2023;41(3):348-354.
16. Anwar S, Jafferany M. Trichotillomania: a psychopathological perspective and the psychiatric comorbidity of hair pulling. 2019.
17. Picardi A, Abeni D, Melchi CF, Puddu P, Pasquini P. Psychiatric morbidity in dermatological outpatients: an issue to be recognized. *Br J Dermatol*. 2000;143(5):983-991.